

(19)



(11)

EP 2 266 573 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
17.06.2015 Bulletin 2015/25

(51) Int Cl.:
A61K 31/565 (2006.01) A61P 35/00 (2006.01)
A61K 47/14 (2006.01) A61K 47/44 (2006.01)
A61K 9/08 (2006.01)

(21) Application number: **10180667.7**

(22) Date of filing: **08.01.2001**

(54) **Fulvestrant formulation**

Fulvestrant Formulierung

Formulation de fulvestrant

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE TR
 Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **10.01.2000 GB 0000313**
12.04.2000 GB 0008837

(43) Date of publication of application:
29.12.2010 Bulletin 2010/52

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
05016921.8 / 1 669 073
01900186.6 / 1 250 138

(73) Proprietor: **AstraZeneca AB**
151 85 Södertälje (SE)

(72) Inventors:
 • **Evans, John**
Macclesfield, Cheshire SK10 4TG (GB)
 • **Grundy, Rosalind Ursula**
Macclesfield, Cheshire SK10 4TG (GB)

(74) Representative: **Hoffmann Eitle**
Patent- und Rechtsanwälte PartmbB
Arabellastraße 30
81925 München (DE)

(56) References cited:
EP-A- 0 310 542 EP-A- 0 346 014
WO-A-96/19997 WO-A-97/21440
US-A- 3 164 520 US-A- 4 388 307

- RIFFKIN C ET AL: "CASTOR OIL AS A VEHICLE FOR PARENTERAL ADMINISTRATION OF STEROID HORMONES.", JOURNAL OF PHARMACEUTICAL SCIENCES AUG 1964, vol. 53, August 1964 (1964-08), pages 891-895, XP009094139, ISSN: 0022-3549
- JOHN C. WATERTON; ET AL.: "A Case of Adenomyosis in a Pigtailed Monkey Diagnosed by Magnetic Resonance Imaging and treated with the Novel Pure Antiestrogen, ICI 182,780", LABORATORY ANIMAL SCIENCE, vol. 43, no. 3, 1993, pages 247-251, XP000998289,
- ROBERTSON J F R ET AL: "Fulvestrant: pharmacokinetics and pharmacology.", BRITISH JOURNAL OF CANCER MAR 2004, vol. 90 Suppl 1, March 2004 (2004-03), pages S7-10, ISSN: 0007-0920
- MCLESKEY S W ET AL: "Tamoxifen-resistant fibroblast growth factor-transfected MCF-7 cells are cross-resistant in vivo to the antiestrogen ICI 182,780 and two aromatase inhibitors.", CLINICAL CANCER RESEARCH : AN OFFICIAL JOURNAL OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH MAR 1998, vol. 4, no. 3, March 1998 (1998-03), pages 697-711, ISSN: 1078-0432
- HOWELL A ET AL: "PHARMACOKINETICS, PHARMACOLOGICAL AND ANTI-TUMOUR EFFECTS OF THE SPECIFIC ANTI-OESTROGEN ICI 182780 IN WOMEN WITH ADVANCED BREAST CANCER", BRITISH JOURNAL OF CANCER, NATURE PUBLISHING GROUP, GB, vol. 74, no. 2, 1 January 1996 (1996-01-01), pages 300-308, XP001007967, ISSN: 0007-0920

EP 2 266 573 B1

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] The invention relates to a novel sustained release pharmaceutical formulation adapted for administration by intra-muscular injection containing the compound 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol in solution in a ricinoleate vehicle which additionally comprises at least one alcohol and a non-aqueous ester solvent which is miscible in the ricinoleate vehicle, for use in the treatment of breast cancer.

[0002] Oestrogen deprivation is fundamental to the treatment of many benign and malignant diseases of the breast and reproductive tract. In premenopausal women, this is achieved by the ablation of ovarian function through surgical, radiotherapeutic, or medical means, and, in postmenopausal women, by the use of aromatase inhibitors.

[0003] An alternative approach to oestrogen withdrawal is to antagonise oestrogens with antioestrogens. These are drugs that bind to and compete for oestrogen receptors (ER) present in the nuclei of oestrogen-responsive tissue. Conventional nonsteroidal antioestrogens, such as tamoxifen, compete efficiently for ER binding but their effectiveness is often limited by the partial agonism they display, which results in an incomplete blockade of oestrogen-mediated activity (Furr and Jordan 1984, May and Westley 1987).

[0004] The potential for nonsteroidal antioestrogens to display agonistic properties prompted the search for novel compounds that would bind ER with high affinity without activating any of the normal transcriptional hormone responses and consequent manifestations of oestrogens. Such molecules would be "pure" antioestrogens, clearly distinguished from tamoxifen-like ligands and capable of eliciting complete ablation of the trophic effects of oestrogens. Such compounds are referred to as Estrogen Receptor-Downregulators (E.R.D.). The rationale for the design and testing of novel, pure antioestrogens has been described in: Bowler et al 1989, Wakeling 1990a, 1990b, 1990c. Wakeling and Bowler 1987, 1988.

[0005] Steroidal analogues of oestradiol, with an alkylsulphonyl side chain in the 7 α position, provided the first examples of compounds devoid of oestrogenic activity (Bowler et al 1989). One of these, 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol was selected for intensive study on the basis of its pure oestrogen antagonist activity and significantly increased antioestrogenic potency over other available antioestrogens. *In vitro* findings and early clinical experience with 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol have promoted interest in the development of the drug as a therapeutic agent for oestrogen-dependent indications such as breast cancer and certain benign gynaecological conditions.

[0006] 7 α -[9-(4,4,5,5,5-Pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, or ICI 182,780, has been allocated the international non-proprietary name fulvestrant, which is used hereinafter. When referring to fulvestrant we include pharmaceutically-acceptable salts thereof and any possible solvates of either thereof.

[0007] Fulvestrant binds to ER with an affinity similar to that of oestradiol and completely blocks the growth stimulatory action of oestradiol on human breast cancer cells *in vitro*; it is more potent and more effective than tamoxifen in this respect. Fulvestrant blocks completely the uterotrophic action of oestradiol in rats, mice and monkeys, and also blocks the uterotrophic activity of tamoxifen.

[0008] Because fulvestrant has none of the oestrogen-like stimulatory activity that is characteristic of clinically available antioestrogens such as tamoxifen or toremifene, it may offer improved therapeutic activity characterised by more rapid, complete, or longer-lasting tumour regression; a lower incidence or rate of development of resistance to treatment; and a reduction of tumour invasiveness.

[0009] In intact adult rats, fulvestrant achieves maximum regression of the uterus at a dose which does not adversely affect bone density or lead to increased gonadotrophin secretion. If also true in humans, these findings could be of extreme importance clinically. Reduced bone density limits the duration of oestrogen-ablative treatment for endometriosis. Fulvestrant does not block hypothalamic ER. Oestrogen ablation also causes or exacerbates hot flushes and other menopausal symptoms; fulvestrant will not cause such effects because it does not cross the blood-brain barrier. **3**

[0010] European Patent Application No. 0 138 504 discloses that certain steroid derivatives are effective antioestrogenic agents. The disclosure includes information relating to the preparation of the steroid derivatives. In particular there is the disclosure within Example 35 of the compound 7 α -[9-(4,4,5,5,5-pentafluoropentylsulphonyl)nonyl]oestra-1,3,5(10)-triene-3,17 β -diol, which compound is specifically named in Claim 4. It is also disclosed that the compounds of that invention may be provided for use in the form of a pharmaceutical composition comprising a steroid derivative of the invention together with a pharmaceutically-acceptable diluent or carrier. It is stated therein that the composition can be in a form suitable for oral or parenteral administration.

[0011] Fulvestrant shows, along with other steroidal based compounds, certain physical properties which make formulation of these compounds difficult. Fulvestrant is a particularly lipophilic molecule, even when compared with other steroidal compounds, and its aqueous solubility is extremely low at around 10 ngml⁻¹ (this is an estimate from a water/solvent mixture solute since measurements this low could not be achieved in a water only solute).

[0012] Currently there are a number of sustained release injectable steroidal formulations which have been commercialised. Commonly these formulations use oil as a solvent and wherein additional excipients may be present. Below in Table 1 are described a few commercialised sustained release injectable formulations.

EP 2 266 573 B1

[0013] In the formulations within Table 1 a number of different oils are used to solubilise the compound and additional excipients such as benzyl benzoate, benzyl alcohol and ethanol have been used. Volumes of oil needed to solubilise the steroid active ingredient are low. Extended release is achievable for periods from 1 to 8 weeks.

5

10

15

20

25

30

35

40

45

50

55

Table 1 - OIL BASED LONG-ACTING INTRAMUSCULAR INJECTIONS

PRODUCT NAME	STEROID	DOSE	TYPE	COMP.	SOURCE	OIL	BzBz	BzOH	EtOH	DOSE	DOSING
SUSTANON 100	Testosterone propionate	30mg	Androgen	Organon	ABPI Data	Arachis		0.1ml		1ml	3 weeks
	Testosterone phenylpropionate	60mg			Sheet Comp.1999						
	Testosterone isocaproate	60mg									
	Testosterone decanoate	100mg									
	Hydroxy progesterone hexanoate	250mgml ⁻¹	Progestogen	Schering HC	ABPI Data Sheet	Castor	up to 46%			1 or 2ml	1 week
TOCOGESTAN	Hydroxy progesterone enantate	200mg	Progestogen	Theramax	Dict. Vidal 1999	Ethyl oleate	*40%			2ml	< 1week
	Progesterone	50mg			Comp.1999						
TROPHOBOLINE	α -Tocopherol	250mg									
	Estrapronicate	1.3mg	Mixed	Theramax	Dict. Vidal 1997	Olive	45%			1ml	15 to 30 days
	Nandrolone undecanoate	50mg									
NORISTERAT	Hydroxyprogesterone heptanoate	80mg									
	Norethisterone oenanthoate	200mg	Contraceptive	Schering HC	ABPI Data Sheet Comp. 1999	Castor	YES			1ml	8 weeks
BENZO-GYNOESTRYL	Estradiol hexahydrobenzoate	5mg	Estradiol	Roussel	Dict. Vidal 1998	Arachis				1ml	1 week
	Hydroxy progesterone caproate	250mgml ⁻¹	Progestogen	Pharlon	Dict. Vidal 1999	Castor	YES			1 or 2ml	1 week
GRAVIBINAN	Estradiol 17- β -valerate	5mgml ⁻¹	Mixed	Schering HC	Dict. Vidal 1995	Castor	YES			1 or 2ml	1 - 2 weeks
	Hydroxyprogesterone caproate	250mgml ⁻¹									
PARABOLAN	Trenbolone	76mg	Androgen	Negma	Dict. Vidal 1997	Arachis		75mg	45mg	1.5ml	2 weeks
DELESTROGEN	Estradiol valerate	20mgml ⁻¹	Estradiol	BMS	J.Pharm. Sci (1964)	Castor	78%	20%	2%		
		40mgml ⁻¹			53(8) 891		58%	40%	2%		

5
10
15
20
25
30
35
40
45
50
55

(continued)

PRODUCT NAME	STEROID	DOSE	TYPE	COMP.	SOURCE	OIL	BzBz	BzOH	EtOH	DOSE	DOSING
DELALUTIN	17-Hydroxy progesterone	250mgml ⁻¹	Progesterogen	DMS	J. Pharm. Sci.(1964) 53(8)891	Castor	YES	YES	up to 2%		

BzBz = benzylbenzoate BzOH = benzylalcohol EtOH = ethanol Dict. Vidal = Dictionnaire Vidal % are w/v and * approximate as measured directly from a single sample

EP 2 266 573 B1

[0014] In US 5,183,814 Example 3 an oil based injection formulation of fulvestrant is described which comprises 50mg of fulvestrant, 400mg of benzyl alcohol and sufficient castor oil to bring the solution to a volume of 1 ml. Manufacture at a commercial scale of a formulation as described in US 5,183,814 will be complicated by the high alcohol concentration. Therefore, there is a need to lower the alcohol concentration in fulvestrant formulations whilst preventing precipitation of fulvestrant from the formulation.

[0015] Table 2 shows the solubility of fulvestrant in a number of different solvents.

Table 2 - SOLUBILITY OF FULVESTRANT

SOLVENT	SOLUBILITY (mgml ⁻¹ at 25°C)
Water	0.001
Arachis oil	0.45
Sesame oil	0.58
Castor oil	20
Miglyol 810	3.06
Miglyol 812	2.72
Ethyl oleate	1.25
Benzyl benzoate	6.15
Isopropyl myristate	0.80
Span 85 (surfactant)	3.79
Ethanol	>200
Benzyl Alcohol	>200

[0016] As can be seen fulvestrant is significantly more soluble in castor oil than any of the other oils tested. The greater solvating ability of castor oil for steroidal compounds is known and is attributed to the high number of hydroxy groups of ricinoleic acid, which is the major constituent of the fatty acids within the triglycerides present in castor oil - see (Riffkin et.al. J. Pharm. Sci., (1964), 53, 891).

[0017] However, even when using the best oil based solvent, castor oil, we have found that it is not possible to dissolve fulvestrant in an oil based solvent alone so as to achieve a high enough concentration to dose a patient in a low volume injection and achieve a therapeutically significant release rate. To achieve a therapeutically significant release rate the amount of fulvestrant needed would require the formulation volume to be large, at least 10 ml. This requires the doctor to inject an excessively large volume of formulation to administer a dose significantly high enough for human therapy.

[0018] Currently guidelines recommend that no more than 5mls of liquid is injected intramuscularly in a single injection. Pharmacologically active doses required for a 1 month long acting depot formulation of fulvestrant is around 250mg. Therefore, when dissolved in just castor oil, fulvestrant would need to be administered in at least 10ml of castor oil.

[0019] The addition of organic solvents in which fulvestrant is freely soluble, and which are miscible with castor oil, may be used, such as an alcohol. With the addition of high concentrations of an alcohol concentrations of >50mgml⁻¹ of fulvestrant in a castor oil formulation is achievable, thereby giving an injection volumes of <5ml - see Table 3 below. We have surprisingly found that the introduction of a non-aqueous ester solvent which is miscible in the castor oil and an alcohol surprisingly eases the solubilisation of fulvestrant into a concentration of at least 50 mgml⁻¹ - see Table 3 below. The finding is surprising since the solubility of fulvestrant in non-aqueous ester solvents - see Table 2 above - is significantly lower than the solubility of fulvestrant in an alcohol. The solubility of fulvestrant is also lower in non-aqueous ester solvents than is the solubility of fulvestrant in castor oil.

[0020] Therefore, we present as a feature of the application a pharmaceutical formulation comprising fulvestrant (preferably fulvestrant is present at 3-10%w/v, 4-9%w/v, 4-8%w/v, 4-7%w/v, 4-6%w/v and most preferably at about 5%w/v) in a ricinoleate vehicle, a pharmaceutically acceptable non-aqueous ester solvent, and a pharmaceutically acceptable alcohol wherein the formulation is adapted for intramuscular administration and attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

[0021] Another feature of the application is a pharmaceutical formulation comprising fulvestrant in which the formulation is adapted for intra-muscular injection into a human and which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

[0022] Further features of the application include a pharmaceutical formulation adapted for intra-muscular injection comprising fulvestrant, 30% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation which is capable after injection of attaining a therapeutically significant blood plasma fulvestrant concentration for at least 2 weeks.

[0023] Further features of the application include a pharmaceutical formulation adapted for intra-muscular injection

EP 2 266 573 B1

comprising fulvestrant; 35% (preferably 30% and ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible within a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.

5 **[0024]** For the avoidance of any doubt when using the term % weight per volume of formulation for the constituents of the formulation we mean that within a unit volume of the formulation a certain percentage of the constituent by weight will be present, for example a 1% weight per volume formulation will contain within a 100ml volume of formulation 1g of the constituent. By way of further illustration

10

% of x by weight per volume of formulation	weight of x in 1ml of formulation
30%	300mg
20%	200mg
10%	100mg
5%	50mg
1%	10mg

15

[0025] Preferred pharmaceutical formulations of the application are as described above wherein:

- 20
1. The total volume of the formulation is 6ml, or less, and the concentration of fulvestrant is at least 45mgml⁻¹.
 2. The total amount of fulvestrant in the formulation is 250mg, or more, and the total volume of the formulation is 6ml, or less.
 3. The total amount of fulvestrant in the formulation is 250mg and the total volume of the formulation is 5-5.25ml.

25 **[0026]** It is appreciated that in the formulation an excess of formulation may be included to allow the attendant physician or care giver to be able to deliver the required dose. Therefore, when a 5ml dose is required it would be appreciated that an excess of up to 0.25ml, preferably up to 0.15ml will also be present in the formulation. Typically the formulation will be presented in a vial or a prefilled syringe, preferably a prefilled syringe, containing a unit dosage of the formulation as described herein, these being further features of the invention.

30 **[0027]** Preferred concentrations of a pharmaceutically-acceptable alcohol present in any of the above formulations are; at least 3%w/v, at least 5%w/v, at least 7%w/v, at least 10% w/v, at least 11% w/v, at least 12% w/v, at least 13% w/v, at least 14% w/v, at least 15% w/v and, preferably, at least 16% w/v. Preferred maximal concentrations of pharmaceutically-acceptable alcohol present in the formulation are ;28% w/v or less, 22% w/v or less and 20% w/v or less.. Preferred ranges of pharmaceutically-acceptable alcohol present in any of the above formulations are selected from any minimum or maximum value described above and preferably are; 3-35%w/v, 4-35%w/v, 5-35%w/v, 5-32%w/v, 7-32%w/v, 10-30%w/v, 12-28%w/v, 15-25%w/v, 17-23%w/v, 18-22%w/v and ideally 19-21%w/v.

35 **[0028]** The pharmaceutically-acceptable alcohol may consist of one alcohol or a mixture of two or more alcohols, preferably a mixture of two alcohols. Preferred pharmaceutically-acceptable alcohols for parenteral administration are ethanol, benzyl alcohol or a mixture of both ethanol and benzyl alcohol, preferably the ethanol and benzyl alcohol are present in the formulation in the same w/v amounts. Preferably the formulation alcohol contains 10% w/v ethanol and 10% w/v benzyl alcohol.

40 **[0029]** The pharmaceutically-acceptable non-aqueous ester solvent may consist of one or a mixture of two or more pharmaceutically-acceptable non-aqueous ester solvents, preferably just one. A preferred pharmaceutically-acceptable non-aqueous ester solvent for parenteral administration is selected from benzyl benzoate, ethyl oleate, isopropyl myristate, isopropyl palmitate or a mixture of any thereof.

45 **[0030]** The ricinoleate vehicle should preferably be present in the formulation in a proportion of at least 30% weight per volume of the formulation, ideally at least 40% or at least 50% weight per volume of formulation.

50 **[0031]** It will be understood by the skilled person that the pharmaceutically-acceptable alcohol will be of a quality such that it will meet pharmacopoeial standards (such as are described in the US, British, European and Japanese pharmacopoeias) and as such will contain some water and possibly other organic solvents, for example ethanol in the US Pharmacopeia contains not less than 94.9% by volume and not more than 96.0% by volume of ethanol when measured at 15.56°C. Dehydrated alcohol in the US Pharmacopeia contains not less than 99.5% ethanol by volume when measured at 15.56°C.

55 **[0032]** Preferred concentrations of the pharmaceutically-acceptable non-aqueous ester solvent present in any of the above formulations are; at least 5% w/v, at least 8% w/v, at least 10% w/v, at least 11% w/v, at least 12% w/v, at least 13% w/v, at least 15% w/v, at least 16% w/v, at least 17% w/v, at least 18% w/v, at least 19% w/v and at least 20% w/v. Preferred maximal concentrations of the pharmaceutically-acceptable non-aqueous ester solvent are; 60% w/v or less,

EP 2 266 573 B1

50%w/v or less, 45% w/v or less, 40% w/v or less, 35% w/v or less, 30% w/v or less and 25% w/v or less. A preferred concentration is 15% w/v. Preferred ranges of pharmaceutically-acceptable non-aqueous ester solvent present in any of the above formulations are selected from any minimum or maximum value described above and preferably are; 5-60%w/v, 7-55%w/v, 8-50%w/v, 10-50%w/v, 10-45%w/v, 10-40%w/v, 10-35%w/v, 10-30%w/v, 10-25%w/v, 12-25%w/v, 12-22%w/v, 12-20%w/v, 12-18%w/v, 13-17%w/v and ideally 14-16%w/v. Preferably the ester solvent is benzyl benzoate, most preferably at about 15%w/v.

[0033] It will be understood by the skilled person that the pharmaceutically-acceptable non-aqueous ester solvent will be of a quality that it will meet pharmacopoeial standards (such as described in the US, British, European and Japanese pharmacopoeias).

[0034] Preferred combinations of pharmaceutically-acceptable alcohol and pharmaceutically-acceptable non-aqueous ester solvent in the formulation are set out below:

Pharmaceutically-acceptable alcohol(%w/v)	Pharmaceutically-acceptable non-aqueous ester (%w/v)
10-30	5-60, 7-55, 8-50, 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 12-25, 12-22, 12-20, 12-18, 13-17 and ideally 14-16.
17-23	5-60, 7-55, 8-50, 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 12-25, 12-22, 12-20, 12-18, 13-17 and ideally 14-16.
3-35, 4-35, 5-35, 5-32, 7-32, 10-30, 12-28, 15-25, 17-23, 18-22 and ideally 19-	10-35
3-35, 4-35, 5-35, 5-32, 7-32, 10-30, 12-28, 15-25, 17-23, 18-22 and ideally 19-21.	12-18
ethanol and benzyl alcohol, most preferably each at about 10%	benzyl benzoate, most preferably at about 15%

[0035] By the use of the term ricinoleate vehicle we mean an oil which has as a proportion (at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 95% w/v) of its composition as triglycerides of ricinoleic acid. The ricinoleate vehicle may be a synthetic oil or conveniently is castor oil, ideally of pharmacopoeial standards, as described above.

[0036] We have surprisingly found that the above formulations of the application provide, after intra-muscular injection, satisfactory release of fulvestrant over an extended period of time.

[0037] This finding is indeed surprising for the following reasons.

1. Previously tested by the applicants have been intra-muscular injections of fulvestrant in the form of an aqueous suspension. We have found extensive local tissue irritation at the injection site as well as a poor release profile. It is believed that the tissue irritation/inflammation was due to the presence of fulvestrant in the form of solid particles. The release profile appeared to be determined by the extent of inflammation/irritation present at the injection site and this was variable and difficult to control. Also the fulvestrant release rate was not sufficiently high to be clinically significant.
2. Our findings from studies using ¹⁴C labelled benzyl alcohol show that it dissipates rapidly from the injection site and is removed from the body within 24 hours of administration.

[0038] It would be expected that ethanol will dissipate at least as quickly, if not more rapidly, from the injection site.

[0039] It is known that benzyl benzoate is metabolised by conjugation to glycine to form hippuric acid by the human liver and excreted into the urine - Martindale: The Extra Pharmacopoeia 32nd edition page 1103, and, therefore, it is unlikely that benzyl benzoate, when used, is present at the injection site during the whole of the extended release period.

[0040] We have found that despite the rapid elimination of the additional solubilising excipients, i.e. the alcohol and pharmaceutically-acceptable non-aqueous ester solvent, from the formulation vehicle and the site of injection after injection of the formulation, extended release at therapeutically significant levels of fulvestrant over an extended period can still be achieved by the formulation of the invention.

[0041] By use of the term "therapeutically significant levels" we mean that blood plasma concentrations of at least 2.5 ngml⁻¹, ideally at least 3 ngml⁻¹, at least 8.5 ngml⁻¹, and up to 12 ngml⁻¹ of fulvestrant are achieved in the patient. Preferably blood plasma levels should be less than 15 ngml⁻¹.

[0042] By use of the term "extended release" we mean at least two weeks, at least three weeks, and, preferably at least four weeks of continuous release of fulvestrant is achieved. In a preferred feature extended release is achieved for 36 days. Preferably extended release of fulvestrant is for at least 2- 5 weeks and more preferably for the following periods (weeks) 2.5-5, 2.5-4, 3-4, 3.5-4 and most preferably for at least about 4 weeks.

[0043] It will be understood that the attendant physician may wish to administer the intramuscular injection as a divided dose, i.e. a 5ml formulation is sequentially administered in two separate injections of 2.5ml, this is a further feature of the invention

[0044] Simply solubilising fulvestrant in an oil based liquid formulation is not predictive of a good release profile or lack of precipitation of drug after injection at the injection site.

[0045] Table 3 shows the solubility of fulvestrant in a castor oil vehicle additionally containing alcohols ethanol and benzyl alcohol with or without benzyl benzoate. The results clearly show the positive effect of benzyl benzoate on fulvestrant solubility in castor oil, despite fulvestrant having a lower solubility in benzyl benzoate than in either alcohol or castor oil.

Table 3

[0046]

Table 3 - EFFECT OF BENZYL BENZOATE ON FULVESTRANT SOLUBILITY IN CASTOR OIL AT 25°C

	% w/v							
Ethanol (96%)	5	5	10	10	10	10	15	15
Benzyl Alcohol	5	5	5	5	10	10	15	15
Benzyl Benzoate		15		15		15		15
Castor Oil	to 100	to 100	to 100	to 100	to 100	to 100	to 100	to 100
Fulvestrant Solubility [mgml ⁻¹]	27	36	46	54	45	65	76	102

[0047] The following Table 4 shows the solubility of fulvestrant in a range of oil based formulations which contain the same amounts of alcohol and benzyl benzoate but in which the oil is changed. The data also shows solubility of fulvestrant after removal of the alcohols.

Table 4

Solubility comparisons of fulvestrant in oil based formulations with and without alcohols

Formulation (a)	Fulvestrant Solubility mg ml ⁻¹ @ 25°C	
	Complete vehicle	Vehicle minus alcohols
Castor oil based	81.2	12.6
Miglyol 812-N based	86.8	1.7
Sesame seed/Castor oil (1:1) based	70.1	4.4
Sesame seed oil based	45.7	0.7
Arachis oil based	40.2	< 0.2

Effect of formulation on precipitation of fulvestrant at the injection site

Formulation ^a	Days						
	2	3	4	7	10	30	51
Formulation F1 castor oil based	0	0	0	0	0	0	0
Formulation F2 Miglyol 812-N based	++ ^b	+++	+++	+++	+++	++	0
Formulation F3 sesame seed oil/castor oil based	+ ^c	++	++	+++	++	+	+

(a) **Complete Vehicle** Formulations comprised ethanol [96%](10%), benzyl alcohol (10%) and benzyl benzoate (15%) made to volume with the stated oil. Excess fulvestrant was added to each solvent mixture and solubility determined. 0, +, ++, +++ = Degree of precipitation (None detected, Mild, Moderate, Severe)
 Formulations comprised fulvestrant (5%), ethanol [96%] (10%), benzyl alcohol (10%) and benzyl benzoate (15%) made to volume with the stated oil.

^b Mainly large needle shaped crystals

^c Small needles and/or sheafs of crystals

[0048] Precipitation of fulvestrant and the release profile was determined with the above formulations in an *in vivo* rabbit study.

[0049] Figure 1 shows the release profile *in vivo* of the four formulations from the second part of Table 4 and shows the effect of the fixed oil component on fulvestrant plasma profile over five days following intramuscular administration in rabbits (data normalised to 50mg per 3kg; mean given; number of animals per timepoint = 8, plasma samples assayed for fulvestrant content using lc-ms/ms detection following solvent extraction). As can be seen the castor oil formulation showed a particularly even release profile with no evidence of precipitation of fulvestrant at the injection site.

[0050] Therefore we present as a further feature of the application an extended release pharmaceutical formulation adapted for intramuscular injection comprising fulvestrant; 35% (preferably 30% or ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation and sufficient amount of a ricinoleate vehicle, taking into account the addition of any further optional pharmaceutically-acceptable excipients, so as to prepare a formulation of at least 45mgml⁻¹ of fulvestrant.

[0051] A feature of the invention is a pharmaceutical formulation adapted for intramuscular injection, as defined above, for use in the treatment of a benign or malignant diseases of the breast or reproductive tract, preferably treating breast cancer, by administration to a human in need of such treatment by intramuscular injection an extended release ricinoleate vehicle based pharmaceutical formulation comprising at least 45mgml⁻¹ of fulvestrant; 35% (preferably 30% or ideally 25%) or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% (preferably at least 5% or ideally 10%) weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible in a ricinoleate vehicle per volume of formulation.

[0052] Preferably 5ml of the intramuscular injection is administered.

[0053] A further feature of the invention is use of fulvestrant in the preparation of a pharmaceutical formulation as describe hereinabove, for the treatment of a benign or malignant disease of the breast or reproductive tract, preferably treating breast cancer.

[0054] Additional excipients commonly used in the formulation field including, for example, an antioxidant preservative, a colorant or a surfactant may be used. A preferred optional excipient is a surfactant.

[0055] As described above fulvestrant is useful in the treatment of oestrogen-dependent indications such as breast cancer and gynaecological conditions, such as endometriosis.

[0056] In addition to fulvestrant another similar type of molecule is currently under clinical investigation. SH-646 (11β-fluoro-7α-(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17β-diol) is also putatively a compound with the same mode of action as fulvestrant and has a very similar chemical structure. It is believed that the compound will also share with fulvestrant similar physical properties and therefore the current invention will also have application with this compound.

[0057] A further feature of the application is a pharmaceutical formulation adapted for intra-muscular injection comprising 11β-fluoro-7α-(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17β-diol; 35% or less weight of a pharmaceutically-acceptable alcohol per volume of formulation, at least 1% weight of a pharmaceutically-acceptable non-aqueous ester solvent miscible within a ricinoleate vehicle per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45mgml⁻¹ of 11β-fluoro-7α-(14,14,15,15,15-pentafluoro-6-methyl-10-thia-6-azapentadecyl)estra-1,3,5(10)-triene-3,17β-diol.

[0058] Further features of the application are those as described above but in which SH-646 is substituted for fulvestrant.

Formulation Example

[0059] Fulvestrant is mixed with alcohol and benzyl alcohol, stirring until completely dissolved. Benzyl benzoate is added and the solution is made to final weight with castor oil and stirred, (for convenience weight is used rather than volume by using the weight to volume ratio). The bulk solution is overlaid with Nitrogen. The solution is sterilised by filtration using one or two filters of 0.2μm porosity. The sterile filtrate is kept under a nitrogen overlay as it is filled under aseptic conditions into washed and depyrogenised, sterile primary containers, for example vials or pre-filled syringes. An overage is included in the primary pack to facilitate removal of the dose volume. The primary packs are overlaid with sterile nitrogen, before aseptically sealing.

See also process flow diagram below

[0060] Quantities of each component of the formulation is chosen according to the required formulation specification, examples are described above. For example quantities are added of each component to prepare a formulation which contains


10% weight per volume of benzyl alcohol

10% weight per volume of ethanol

15% weight per volume of benzyl benzoate
 250mg of fulvestrant for each 5ml of finished formulation
 and the remaining amount as castor oil

5

FLOW DIAGRAM OF MANUFACTURING



10

15

20

25

30

35

40

References

45

[0061]

50

55

1. Bowler J, Lilley TJ, Pittam JD, Wakeling AE. Novel steroidal pure antioestrogens. Steroids 989; 5471-99.
2. Wakeling AE. Novel pure antioestrogens: mode of action and therapeutic prospects. American New York Academy Science 1990a; 595: 348-56.
3. Wakeling AE. Steroidal pure antioestrogens. In Lippman M, Dickson R, editors. Regulatory mechanisms in breast cancer. Boston: Kluwer Academic, 1990b: 239-57.
4. Wakeling AE. Therapeutic potential of pure antioestrogens in the treatment of breast cancer. Journal Steroid Biochemistry 1990c; 37: 771-5.
5. Wakeling AE, Bowler J. Steroidal pure antioestrogens. Journal Endocrinology 1987; 112: R7-10.

6. Wakeling AE, Bowler J. Biology and mode of action of pure antioestrogens. Journal Steroid Biochemistry 1988; 3: 141-7.

5 **Claims**

- 10 1. A pharmaceutical formulation for use in the treatment of breast cancer by intra-muscular injection, wherein the pharmaceutical formulation comprises fulvestrant, a pharmaceutically-acceptable alcohol being a mixture of 10 % weight of ethanol per volume of formulation and 10 % weight of benzyl alcohol per volume of formulation, and the formulation contains 15 % weight of benzyl benzoate per volume of formulation and a sufficient amount of a ricinoleate vehicle so as to prepare a formulation of at least 45 mgml⁻¹ of fulvestrant, wherein the ricinoleate vehicle is castor oil, and wherein the total volume of the formulation is 6 ml or less.
- 15 2. The pharmaceutical formulation as claimed in claim 1 wherein the total amount of fulvestrant in the formulation is 250 mg, or more, and the total volume of the formulation is 6 ml, or less.
3. A pharmaceutical formulation as claimed in claim 2 wherein the total amount of fulvestrant in the formulation is 250 mg and the total volume of the formulation is 5 to 5.25 ml.

20

Patentansprüche

- 25 1. Pharmazeutische Formulierung zur Verwendung bei der Behandlung von Brustkrebs mittels intramuskulärer Injektion, wobei die pharmazeutische Formulierung Fulvestrant und einen pharmazeutisch annehmbaren Alkohol umfasst, der ein Gemisch aus 10 Gew.-% Ethanol, bezogen auf das Volumen der Formulierung, und 10 Gew.-% Benzylalkohol, bezogen auf das Volumen der Formulierung, darstellt, und wobei die Formulierung 15 Gew.-% Benzylbenzoat, bezogen auf das Volumen der Formulierung, und eine zur Herstellung einer Formulierung mit mindestens 45 mgml⁻¹ Fulvestrant ausreichende Menge einer Ricinoleat-Trägersubstanz enthält, wobei die Ricinoleat-Trägersubstanz Rizinusöl ist, und wobei das Gesamtvolumen der Formulierung 6 ml oder weniger ist.
- 30 2. Die pharmazeutische Formulierung wie in Anspruch 1 beansprucht, wobei die Gesamtmenge an Fulvestrant in der Formulierung 250 mg oder mehr und das Gesamtvolumen der Formulierung 6 ml oder weniger ist.
- 35 3. Die pharmazeutische Formulierung wie in Anspruch 2 beansprucht, wobei die Gesamtmenge an Fulvestrant in der Formulierung 250 mg und das Gesamtvolumen der Formulierung 5 bis 5,25 ml ist.

Revendications

- 40 1. Formulation pharmaceutique pour utilisation dans le traitement du cancer du sein par injection intramusculaire, la formulation pharmaceutique comprenant du fulvestrant, un alcool pharmaceutiquement acceptable étant un mélange de 10 % en poids d'éthanol par volume de formulation et 10 % en poids d'alcool benzylique par volume de formulation, et la formulation contient 15 % en poids de benzoate de benzyle par volume de formulation et une quantité suffisante d'un véhicule de ricinoléate de manière à préparer une formulation d'au moins 45 mg.ml⁻¹ de fulvestrant, le volume total de la formulation étant de 6 ml ou moins.
- 45 2. Formulation pharmaceutique selon la revendication 1 dans laquelle la quantité totale de fulvestrant dans la formulation est de 250 mg, ou plus, et le volume total de la formulation est de 6 ml, ou moins.
- 50 3. Formulation pharmaceutique selon la revendication 2 dans laquelle la quantité totale de fulvestrant dans la formulation est de 250 mg, et le volume total de la formulation est de 5 à 5,25 ml.

55




Figure 1

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- EP 0138504 A [0010]
- US 5183814 A [0014]

Non-patent literature cited in the description

- RIFFKIN. *J. Pharm. Sci.*, 1964, vol. 53, 891 [0016]
- BOWLER J ; LILLEY TJ ; PITTAM JD ; WAKELING AE. Novel steroidal pure antioestrogens. *Steroids*, vol. 989, 5471-99 [0061]
- WAKELING AE. Novel pure antioestrogens: mode of action and therapeutic prospects. American New York Academy Science, 1990, vol. 595, 348-56 [0061]
- Steroidal pure antioestrogens. WAKELING AE. Regulatory mechanisms in breast cancer. Kluwer Academic, 1990, 239-57 [0061]
- WAKELING AE. Therapeutic potential of pure antioestrogens in the treatment of breast cancer. *Journal Steroid Biochemistry*, 1990, vol. 37, 771-5 [0061]
- WAKELING AE ; BOWLER J. Steroidal pure antioestrogens. *Journal Endocrinology*, 1987, vol. 112, R7-10 [0061]
- WAKELING AE ; BOWLER J. Biology and mode of action of pure antioestrogens. *Journal Steroid Biochemistry*, 1988, vol. 3, 141-7 [0061]